

## Zooplankton distribution in coastal water of the North-Western Bay of Bengal, off Rushikulya estuary, east coast of India

\*S. Srichandan, Biraja K. Sahu, R. Panda, S. K. Baliarsingh<sup>#</sup>, K. C. Sahu & R. C. Panigrahy

Department of Marine Sciences, Berhampur University, Berhampur-760007, Odisha, India

<sup>#</sup>(Present address) Indian National Centre for Ocean Information Services, Hyderabad-500090, India

\*[E-mail: [suchismita.sima@gmail.com](mailto:suchismita.sima@gmail.com)]

Received 17 September 2013; revised 6 June 2014

### Abstract

In total, 186 species of holoplankton and 23 different types of meroplankton were encountered during this study. Zooplankton community was dominated by copepods represented by 112 species, belonging to 4 orders and 26 families. Calanoida emerged as the most dominant order among the copepods being comprised of 68 species. It was followed by poecilostomatoids with 26 species, cyclopoids with 10 species and harpacticoida with 8 species. *Paracalanus aculeatus* emerged as the most dominant species during pre-monsoon season while, *Acrocalanus longicornis* was dominant in monsoon and *A. gibber* in the post-monsoon season. Other dominant copepods were *Oithona* sp., *Miracia efferata*, *Acartia southwelli*, *Centropages tenuiremis*, *Paracalanus parvus*, *Acrocalanus gracilis* and *Acartia erythraea*. Average zooplankton density ranged from 2387 org./10m<sup>3</sup> to 11659 org./10m<sup>3</sup>. Zooplankton volume ranged from 0.65 ml/10m<sup>3</sup> in monsoon to 1.51 ml/10m<sup>3</sup> in post-monsoon season. Despite high species abundance during premonsoon period, species diversity was maximum in monsoon. Species richness and dominance indices remained higher during post-monsoon whereas Pielou's evenness (J') was more in monsoon. Non-metric multidimensional scaling (MDS) ordinations based on Bray-Curtis similarities indicated that species composition was unequal during different months as there was no similarity above 40% level.

[**Keywords:** Zooplankton, Copepod, Coastal water, Hydrography, Bay of Bengal]

### Introduction

Zooplankton constitutes a broad category and wide range of organisms in marine environment. Roman *et al*<sup>1</sup> have opined that zooplankton are very important to marine pelagic ecosystems, that support higher trophic levels and as the essential determinant of the potential fishery yield. Several microzooplankton species also constitute major food stuff of the larvae of crustaceans, molluscs and fishes while some species of zooplankton are used as water quality indicators<sup>2</sup> and movement of water current<sup>3</sup>.

Therefore knowledge on species composition, abundance and distribution of zooplankton was always considered as great significance in marine ecological and fishery management exercises<sup>4</sup>. In recognition of the ecological and economic significance of marine zooplankton, emphasis has been laid to acquire more and more knowledge on species composition, seasonal abundance and reproductive biology of marine and estuarine plankton worldwide leading to accumulation of a plethora of literature. Information relating to zooplankton of coastal waters and estuaries in

India has started in early 1900s<sup>5</sup> and it gained momentum from 1950s and 1960s especially after the IIOE. However, majority of studies were confined to areas like Cochin Back waters<sup>6</sup>, Mandovi and Zuari estuaries and their neighboring sea<sup>7</sup>, Vellar estuary and its adjoining coastal water<sup>8</sup> and Hooghly estuary<sup>9</sup>. Zooplankton studies along Odisha coast in general remained meager and were limited to the Chilika lake<sup>10-12</sup>, Rushikulya estuary<sup>13</sup>, Bahuda estuary<sup>14</sup>, Burhabalanga estuary<sup>15</sup>, Mahanadi estuary<sup>16</sup>. The only study on zooplankton distribution in coastal waters along the Odisha coast was that of Sahu *et al*<sup>17</sup>. Present paper describes the seasonal variations in zooplankton species composition, population density and relative abundance of major groups in relation to variations in the physico-chemical parameters of north-western Bay of Bengal, off the Rushikulya estuary.

### Materials and Methods

The present study was carried out in the Bay of Bengal, off Rushikulya estuary, south Odisha coast. Samples were collected from five stations (Fig.1) at monthly interval from March 2010 to

February 2011. This study area is influenced by fresh water flow via Rushikulya river which receives high quantity of dissolved chemical inputs from many sources including the effluent discharges of a Chloro-alkali plant. The physiographic features, climate, tidal rhythms of the region etc have been discussed earlier<sup>13,18</sup>.

Surface water samples were collected using a clean plastic bucket for measurement of hydrographic parameters, nutrients and chlorophyll *a*. Water temperature (WT) and pH were recorded using mercury filled centigrade thermometer and EUTECH field pH reader (accuracy  $\pm 0.01$ ) respectively. Water samples for




Fig.1. Map showing the sampling locations off Rushikulya estuary

analysis of salinity, nutrients and chlorophyll *a* were collected in clean polyethylene bottles and transported to the laboratory kept in an ice box. Samples for dissolved oxygen (DO) analysis were fixed onboard with Winkler's A and B solutions and were transported to laboratory for further analysis. DO was measured adopting Winkler's method, while salinity was estimated following Knudsen's method<sup>19</sup>. Water samples for nutrient analysis were filtered against glass fiber filter (GF/C) using Millipore Filtering System and were analysed for nitrite ( $\text{NO}_2\text{-N}$ ), nitrate ( $\text{NO}_3\text{-N}$ ), ammonia ( $\text{NH}_4\text{-N}$ ), inorganic phosphate ( $\text{PO}_4\text{-P}$ ) and silicate ( $\text{SiO}_4\text{-Si}$ ) following the procedures described by Grasshoff *et al.*<sup>19</sup>. Chlorophyll *a* was

estimated adopting the spectrophotometric method given by Parsons *et al.*<sup>20</sup>.

Zooplankton samples were collected at each station by horizontal haul using a zooplankton net (mesh size of 120  $\mu\text{m}$ ). A digital flow meter (Hydro Bios) was used to determine the volume of water filtered. Samples collected from the receiver were transferred to pre-cleaned polythene bottles and preserved with 5% formaldehyde. In the laboratory, the zooplankton samples were first used for determination of plankton volume through volume displacement method. The residual plankton on the filter paper was re-suspended in 5% formaldehyde and sub-sampling was made using Folsom plankton divider. Subsample was used for quantitative and qualitative analysis. Larger organisms like copepods of families Pontellidae, Eucalanidae, Euchaetidae, Chaetognaths, Dollioloids & Salps were sorted out and counted. Remaining mixture was made to exactly 100 ml and 1 ml of aliquot was transferred on to a Sedgwick-Rafter cell and observed under an inverted microscope (Cippon; Model No.21033) for identification and counting. Total number of macro and micro-zooplankton was taken as the population density. The numerical abundance was presented as  $\text{org./10m}^3$ . Relative abundance was computed from the total density and the density of each group. Identification of zooplankton was done following standard literature<sup>21-23</sup>.

Data were grouped into three seasons: premonsoon (PRM), monsoon (MON) and post-monsoon (POM). Analysis of Variance (ANOVA) test was performed to know the inter season comparison of different parameters. Correlation analysis was made to find out the possible relationships between different parameters and zooplankton density. Species diversity indices i.e. Marglef's Species Richness (*d*), Pielou's Evenness (*J'*), Shannon Wiener Diversity Index (*H'* (log<sub>e</sub>)) and Simpson's Dominance Index ( $\lambda$ ) were determined using PRIMER (Plymouth Routines in Multivariate Ecological Research) software Version 5. Zooplankton species composition and abundance from March 2010 - February 2011 was analysed using nonparametric multivariate statistical methods in PRIMER Version 5<sup>24,25</sup> to decipher the similarity in species composition.

## Results and Discussion

Seasonally pooled mean and standard deviation values of the hydrographic parameters and nutrients are given in Table 1. Each of the

hydrographic parameter such as water temperature (WT), salinity, pH and DO showed well marked seasonal and spatial variations. Monthly collected data showed that water temperature varied from 21.8°C (station R-1, Jan) to 31.4°C (station R-4, May). Seasonally pooled mean temperature values however ranged from 26.0°C to 28.9°C. Higher temperature was recorded in PRM season while lower temperature was observed during POM months. Temperature

over the Bay of Bengal mainly depends on climatic variations especially the atmospheric conditions and rain fall<sup>26</sup>. Temperature variations during the present study could also be ascribed to seasonal changes in atmospheric conditions and rainfall. Monthly recorded salinity values ranged from 25.3 PSU (station R-1, August) to 35.8 PSU (station R-5, January). Seasonally pooled average salinity ranged from 30.58 PSU during MON season to 31.79 PSU during PRM period.

Table 1. Variations of physicochemical and biological parameters

St.	WT (°C)	pH	DO (mg/L)	Salinity (PSU)	NO <sub>2</sub> (µmol/L)	NO <sub>3</sub> (µmol/L)	NH <sub>4</sub> (µmol/L)	PO <sub>4</sub> (µmol/L)	SiO <sub>4</sub> (µmol/L)	Chl- <i>a</i> (mg/ m <sup>3</sup> )	ZooBio (ml/10m <sup>3</sup> )	ZooPop (Org./10m <sup>3</sup> )
<b>PREMONSOON</b>												
R-1	27.7	7.66	8.02	28.89	0.36	2.06	2.4	2.14	6.13	5.67	0.98	23054
R-2	28.3	7.97	7.78	29.7	0.32	1.66	2.47	1.25	4.72	2.51	1.39	11556
R-3	29.0	8.24	7.64	32.27	0.4	1.4	2.15	1.16	5.32	2.59	0.80	11234
R-4	29.6	8.29	7.86	33.26	0.27	0.93	1.37	0.91	5.02	1.59	1.37	8184
R-5	30.0	8.02	7.67	34.83	0.4	1.08	1.47	1.59	4.58	1.7	0.14	4270
<b>Mean</b>	28.9	8.04	7.79	31.79	0.35	1.43	1.97	1.41	5.15	2.81	0.93	11659
<b>SED</b>	1.0	0.25	0.15	2.47	0.06	0.45	0.52	0.48	0.62	1.66	0.51	7012
<b>MONSOON</b>												
R-1	26.3	7.51	8.02	26.94	0.46	1.26	1.72	2.41	7	2.16	1.38	5958
R-2	26.7	7.77	7.71	28.91	0.49	1.17	2.07	1.46	5.45	1.81	0.86	2416
R-3	27.1	8.07	7.64	32.31	0.42	0.93	1.35	1.31	4.82	1.99	0.54	1077
R-4	27.9	8.27	7.43	31.43	0.37	0.98	1.81	0.77	5.83	1.48	0.35	1129
R-5	27.9	8.3	7.31	33.33	0.3	0.97	1.67	0.85	3.11	1.29	0.16	1354
<b>Mean</b>	27.2	7.98	7.62	30.58	0.41	1.06	1.72	1.36	5.24	1.75	0.65	2387
<b>SED</b>	0.7	0.34	0.27	2.61	0.08	0.14	0.26	0.66	1.43	0.36	0.48	2069
<b>POST MONSOON</b>												
R-1	24.7	8.11	8.46	27.66	0.38	1.82	2.19	2.98	9.23	1.75	2.53	10226
R-2	26.0	8.14	7.32	30.08	0.29	2.05	2.24	1.29	7.75	1.73	1.55	4500
R-3	26.2	8.15	7.43	30.95	0.42	1.89	5.06	1.01	6.01	1.85	1.59	8848
R-4	26.4	8.18	7.58	32.54	0.42	1.28	1.8	1.04	6.78	1.84	0.33	2619
R-5	26.5	8.19	7.29	32.62	0.28	1.24	2.53	0.95	7.07	1.85	1.52	2433
<b>Mean</b>	26.0	8.15	7.62	30.77	0.36	1.66	2.76	1.45	7.37	1.8	1.51	5725
<b>SED</b>	0.7	0.03	0.49	2.05	0.07	0.37	1.31	0.86	1.21	0.06	0.78	3606

Trend of salinity variation was PRM >POM >MON. Lower salinity values during MON season could be attributed to the combined influence of rainfall and freshwater influx from landward sources. Similar type of seasonal variation in salinity was observed in coastal waters of Kalpakkam, East coast of India by Satpathy *et al*<sup>27</sup>. Temperature showed positive correlation ( $p < 0.05$ ) with salinity indicating their strong affinity. Results of one way ANOVA (Table 2) also explained well defined seasonal variation in water temperature ( $p < 0.01$ ). Monthly measured pH values varied between 7.14 (station R-1, July) and 8.58 (station R-4, September), whereas the seasonally pooled mean value lied

between 7.98 (MON) and 8.15 (POM). Average pH in PRM season was 8.04. The higher values of POM and PRM seasons could be due to high salinity and high photosynthetic activity as observed by Subramanian and Mahadevan<sup>28</sup> along Chennai coast. Monthly observed DO values ranged from 6.45 (station R-4, Dec.) – 8.75 (station R-4, Nov). Season-wise average DO values showed that maximum value of 7.79 mg /L was encountered in the PRM period. More fluctuation in DO was however observed during POM season (SD: 0.49 mg /L). High surface values of DO during PRM season could be ascribed to the addition of DO by phytoplankton

photosynthesis as reported in Kalpakkam coastal waters<sup>29</sup>.

Table 2. One way ANOVA results of hydrographic and biological parameters 2010-11

Parameters	p Value
WT	<b>0.000</b>
pH	0.273
DO	0.461
Salinity	0.304
NO <sub>2</sub>	0.484
NO <sub>3</sub>	<b>0.002</b>
NH <sub>4</sub>	0.193
PO <sub>4</sub>	0.976
SiO <sub>4</sub>	<b>0.022</b>
Chl- <i>a</i>	<b>0.042</b>
Zoo Density	<b>0.005</b>

\*Significant level  $p < 0.05$  in bold

Nitrite (NO<sub>2</sub>-N), the most unstable form of nitrogen in sea water, exhibited wide range of variation. The monthly values varied from 0.06  $\mu\text{mol/L}$  (station R-5, Feb) to 0.85  $\mu\text{mol/L}$  (station R-4, Nov). Season-wise pooled mean NO<sub>2</sub>-N was maximum during MON (0.41  $\mu\text{mol/L}$ ), while it remained minimum during the PRM (0.35  $\mu\text{mol/L}$ ) season. Monthly values of NO<sub>3</sub>-N ranged from 0.58 (station R-4, Feb) to 3.13  $\mu\text{mol/L}$  (station R-3, Feb). The negative correlation ( $r = -0.552$ ,  $p < 0.05$ ) between salinity and nitrate denotes that freshwater forms major contributor of nitrate in coastal waters. Monthly NH<sub>4</sub>-N varied from 0.8 (station R-5, Feb) to 15.4  $\mu\text{mol/L}$  (station R-3, Dec) while seasonal average values ranged between 1.72 (MON) to 2.76 (POM)  $\mu\text{mol/L}$ . Monthly values of PO<sub>4</sub>-P varied between 0.17 - 8.86  $\mu\text{mol/L}$  with maximum concentration observed at station R-1 in November 2010. Average seasonal values however ranged from 1.36 (MON) to 1.45 (POM)  $\mu\text{mol/L}$ . Like nitrate, phosphate concentrations also exhibited negative correlation with salinity ( $r = -0.712$ ,  $p < 0.01$ ), indicating thereby this nutrient too is controlled by fresh water influx. The SiO<sub>4</sub>-Si concentration ranged between 0.64 (station R-5, June) to 18.42 (station R-2, Feb)  $\mu\text{mol/L}$ . SiO<sub>4</sub>-Si concentration was maximum ( $7.37 \pm 1.21$   $\mu\text{mol/L}$ ) in POM as compared to PRM ( $5.15 \pm 0.62$   $\mu\text{mol/L}$ ) and MON ( $5.24 \pm 1.43$   $\mu\text{mol/L}$ ) seasons. The strong negative correlation ( $r = -0.583$ ,  $p < 0.05$ ) of silicate and salinity suggests its input via freshwater. Results of ANOVA test (Table 2) showed more

significant seasonal variation in NO<sub>3</sub>-N ( $p < 0.01$ ) and SiO<sub>4</sub>-Si ( $p < 0.05$ ) contents in the study area. The higher value of nutrients in MON season and their negative correlation with salinity establishes their entry through freshwater as was reported along the Coromandel Coast<sup>30</sup> and coastal waters of Gopalpur<sup>31</sup>.

Phytoplankton is the ultimate source of food for zooplankton. Chlorophyll *a* is the best index of phytoplankton standing stock as such it has direct bearing on the growth of zooplankton. Seasonally pooled mean chlorophyll *a* was at its maximum in PRM (2.81  $\text{mg/m}^3$ ) that fell to minimum in MON (1.75  $\text{mg/m}^3$ ). Higher chlorophyll *a* contents in PRM season is a common feature in surface waters of the Bay of Bengal. When all the seasons were taken in to consideration for ANOVA, significant seasonal variation was observed in chlorophyll *a* ( $p < 0.05$ ) distribution.

Monthly values of plankton volume ranged from 0.02 (station R-4, May and station R-3 June) to 6.40  $\text{ml}/10\text{m}^3$  (station R-1, Feb). Higher volume of plankton at station R-1 during Feb might be due to the presence of caridean larvae, brachyuran zoea larvae, chaetognaths and larger sized copepod species. Seasonally pooled mean plankton volume ranged from 0.65  $\text{ml}/10\text{m}^3$  in MON to 1.51  $\text{ml}/10\text{m}^3$  in POM. In PRM, the pooled mean of plankton volume was 0.93  $\text{ml}/10\text{m}^3$ . More fluctuation (SD: 0.78  $\text{ml}/10\text{m}^3$ ) in plankton volume was observed during POM season followed by pre-monsoon (SD: 0.51  $\text{ml}/10\text{m}^3$ ) and monsoon (SD: 0.48  $\text{ml}/10\text{m}^3$ ). Significant positive correlation ( $r = 0.275$ ,  $p < 0.05$ ) was noticed between zooplankton volume and density.

The systematic account of zooplankton encountered during the course of study are presented in Table 3 and Table 4. In total, 186 species of holoplankton belonging to 12 groups such as Acantharia, Ciliophora, Foraminifera, Hydrozoa, Ctenophora, Gastropoda, Cladocera, Copepoda, Ostracoda, Malacostraca, Chaetognatha, Chordata and 23 different types of meroplankton were recorded. The holoplankton components depending upon their abundance are broadly described under three groups namely copepods, other crustaceans and non crustaceans. Copepods formed the most dominant group throughout the study period.

Copepod not only remained as the most dominant group but also appeared in large numbers at all the stations round the year. They were represented by 112 species, belonging to 26 families and 4 orders. Calanoida emerged as the

most dominant group comprising 68 species followed by poecilostomatoids with 26 species, cyclopoids with 10 species and harpacticoid with 8 species. The dominance of calanoida over others could be due to their continuous breeding, quick larval development and adaptation to wide range of environmental conditions as reported earlier<sup>8,32</sup>. Further, among the calanoida, *Paracalanus aculeatus* remained as the most dominant component in PRM season, while *A. longicornis* remained dominant in MON and *A. gibber* in POM season. Rakesh *et al*<sup>33</sup> have reported that both *Paracalanus* sp. and *Acrocalanus* sp. were characterizing the coastal locations off North Coastal Andhra Pradesh and the findings of the

present study corroborated the same. *Canthocalanus pauper* and *Acrocalanus longicornis* were frequently observed throughout the year justifying their cosmopolitan behavior. Among cyclopoida, *Oithona similis*, *O. brevicornis*, *O. spirostris*, *Oithona* sp. appeared as common species in all the three seasons. High abundance of *Oithona* sp. could be due to its high reproductive capability as opined by Santhanam and Perumal<sup>8</sup> and Santosh Kumar and Perumal<sup>34</sup>. Although species like *Miracia efferata*, *Macrosetella gracilis*, *Macrosetella oculata*, *Microsetella norvegica*, *Microsetella rosea*,

Table 3. Check list of zooplankton (Holoplankton) encountered in the coastal waters of south Odisha coast, off Rushikulya estuary during 2010-11

Phylum:Class	S.class	Order	Family	Species
Protozoa:				
Acantharia		Arthracanthida	Acanthometridae	<i>Acanthometron</i> sp. J. Muller (1856)
Ciliophora:				
Spirotrichea	Choreotrichia	Tintinnida	Tintinnidiidae	<i>Leprotintinnus nordqvisti</i> Brandt (1906)
			Codonellidae	<i>Tintinnopsis beroidea</i> Stein (1867)
				<i>T. butschlii</i> Daday (1887)
				<i>T. cylindrica</i> Daday (1887)
				<i>T. mortensenii</i> Schmidt (1901)
				<i>T. tocantinensis</i> Kofoid and Campbell (1929)
				<i>T. tubulosa</i> Levander (1900)
				<i>T. uruguayensis</i> Balech(1948)
			Codonellopsidae	<i>Codonellopsis ostenfeldi</i> Schmidt (1901)
			Dictyocystidae	<i>Dictyocysta seshaiyai</i> Krishnamurthy & Santhanam (1975)
			Metacyclididae	<i>Metacyclis jorgenseni</i> Cleve (1902)
			Rhabdonellidae	<i>Rhabdonella</i> sp. Brandt (1906)
			Tintinnidae	<i>Amphorellopsis</i> sp. Kofoid and Campbell (1929)
				<i>Eutintinnus tenue</i> Kofoid and Campbell (1929)
			Xystonellidae	<i>Favella philippinensis</i> Roxas (1941)
Foraminifera:				
Polythalamia		Globigerinida	Globigerinidae	<i>Globigerina bulloides</i> d'Orbigny (1826)
				<i>G. rubescense</i> Hofker (1956)
				<i>Globigerina</i> sp. d'Orbigny (1826)
		Rotaliida	Rotaliidae	<i>Asterorotalia trispinosa</i> Thalmann (1933)
		Miliolida	Miliolidae	<i>Quinqueloculina</i> sp. d'Orbigny(1826)
Cnidaria:				
Hydrozoa	Hydroidomedusae	Anthomedusae	Cladonemidae	<i>Cladonema</i> sp. Dujardin (1843)
			Corynidae	<i>Sarsia</i> sp. Lesson(1843)
			Hydractiniidae	<i>Podocoryne</i> sp. Luetken (1850)
		Leptomedusae	Campanulariidae	<i>Obelia</i> sp. Péron & Lesueur (1810)
			Phialellidae	<i>Phialella quadrata</i> Forbes (1848)
		Trachymedusae	Geryoniidae	<i>Liriope tetraphylla</i> Chamisso and Eysenhardt (1821)
			Rhopalonematidae	<i>Aglaura hemistoma</i> Peron and Lesueur (1810)
		Conica	Aequoreidae	<i>Aequorea vitrina</i> Gosse (1853)
	Siphonophorae	Physonectae	Agalmatidae	<i>Agalma elegans</i> Sars (1846)

		Calycephorae	Diphyidae	<i>Sulculeolaria</i> sp. Blainville (1830) <i>Diphyes chamissonis</i> Huxley (1859) <i>D. dispar</i> Chamisso and Eysenhardt (1821) <i>Diphyes</i> sp. Cuvier (1817) <i>Lensia</i> sp Totton (1932) <i>Eudoxoides mitra</i> Huxley (1859) <i>Muggiaea</i> sp. Busch (1851)
			Abylidae	<i>Abylopsis</i> sp. Chun (1888) <i>Bassia bassensis</i> Quoy and Gaimard (1833) 1834)
Ctenophora: Tentaculata Ctenophora: Nuda Mollusca: Gastropoda	Typhlocoela	Cydippida	Pleurobrachiidae	<i>Pleurobrachia pileus</i> O. F. Müller (1776)
		Beroida	Beroidae	<i>Beroe</i> sp. Gronov (1760)
	Streptoneura	Mesogastropoda	Atlantidae	<i>Atlanta</i> sp. Lesueur (1817)
			Janthinidae	<i>Janthina</i> sp. Röding (1798)
	Euthyneura	Thecosomata	Limacinidae	<i>Limacina bulimoides</i> d'Orbigny (1834) <i>Limacina inflata</i> d'Orbigny (1834)
			Cavoliniidae	<i>Cresis acicula</i> Rang (1828) <i>Hyalocylix striata</i> Rang (1828)
Arthropoda: Branchiopoda	Phyllopoda	Cladocera	Podonidae	<i>Evadne tergestina</i> Claus (1864)
			Sididae	<i>Penilia avirostris</i> Dana (1849)
Arthropoda: Maxillipoda	Copepoda	Calanoida	Metridinidae	<i>Pleuromamma</i> sp. Giesbrecht in Giesbrecht & Schmeil (1898)
			Acartiidae	<i>Acartia . centrura</i> Giesbrecht (1889) <i>A. danae</i> Giesbrecht (1889) <i>A. erythraea</i> Giesbrecht (1889) <i>A. spinicauda</i> Giesbrecht (1889) <i>A. negligens</i> Dana (1849) <i>A. southwelli</i> Sewell (1914) <i>Acartia</i> sp. Dana (1846)
			Candaciidae	<i>Candacia catula</i> Giesbrecht (1889) <i>Candacia</i> sp. Dana (1846) <i>Paracandacia truncata</i> Dana (1849) <i>Paracandacia</i> sp. Grice (1963)
			Centropagidae	<i>Centropages alcocki</i> Sewell (1912) <i>C. furcatus</i> Dana (1849) <i>C. calaninus</i> Dana (1849) <i>C. dorsispinatus</i> Thompson I.C. & Scott A.(1903) <i>C. orsinii</i> Giesbrecht (1889) <i>C. tenuiremis</i> Thompson I.C. & Scott A.(1903) <i>Centropages</i> sp. Kroyer (1849)
			Pontellidae	<i>Calanopia minor</i> A. Scott (1902) <i>C. elliptica</i> Dana (1846, 1849) <i>Labidocera acuta</i> Dana (1849) <i>L. detruncata</i> Dana (1849) <i>L. minuta</i> Giesbrecht (1889) <i>L. pectinata</i> Thompson I.C. & Scott A. (1903)

---

		<i>Labidocera</i> sp. Lubbock (1853)
		<i>Pontella fera</i> Dana (1849)
		<i>P. securifer</i> Brady (1883)
		<i>Pontellina platychela</i> Fleminger and Hülsemann (1974)
	Temoridae	<i>Temora discaudata</i> Giesbrecht (1889)
		<i>T. turbinata</i> Dana (1849)
		<i>T. stylifera</i> Dana (1849)
	Tortanidae	<i>Tortanus barbatus</i> Brady (1883)
		<i>T. forcipatus</i> Giesbrecht (1889)
		<i>T. gracilis</i> Brady (1883)
	Calanidae	<i>Mesocalanus tenuicornis</i> Dana (1849)
		<i>Canthocalanus pauper</i> Giesbrecht (1888)
		<i>Nannocalanus minor</i> Claus (1863)
		<i>Undinula vulgaris</i> Dana (1849)
	Paracalanidae	<i>Acrocalanus gracilis</i> Giesbrecht (1888)
		<i>A. longicornis</i> Giesbrecht (1888)
		<i>A. gibber</i> Giesbrecht (1888)
		<i>Acrocalanus</i> sp. Giesbrecht (1888)
		<i>Calocalanus pavo</i> Dana (1849)
		<i>Paracalanus aculeatus</i> Giesbrecht (1888)
		<i>P. parvus</i> Claus (1863)
		<i>Paracalanus</i> sp. Boeck (1865)
	Eucalanidae	<i>Eucalanus attenuatus</i> Dana (1849)
		<i>E. monachus</i> Giesbrecht (1888)
		<i>Eucalanus</i> sp. Dana (1852)
		<i>Subeucalanus crassus</i> Giesbrecht (1888)
		<i>S. mucronatus</i> Giesbrecht (1888)
		<i>S. pileatus</i> Giesbrecht (1888)
		<i>S. subcrassus</i> Giesbrecht (1888)
		<i>S. subtenuis</i> Giesbrecht (1888)
	Aetideidae	<i>Undeuchaeta</i> sp. Giesbrecht (1888)
	Arietellidae	<i>Metacalanus aurivilli</i> Cleve (1901)
	Clausocalanidae	<i>Clausocalanus arcuicornis</i> Dana (1849)
	Euchaetidae	<i>Euchaeta concinna</i> Dana (1849)
		<i>E. marina</i> Prestandrea (1833)
		<i>Euchaeta</i> sp. Philippi (1843)
	Scolecitrichidae	<i>Scolecithrix danae</i> Lubbock (1856)
		<i>Scolecithricella minor</i> Brady (1883)
	Pseudodiaptomidae	<i>Pseudodiaptomus aurivilli</i> Cleve (1901)
		<i>P. serricaudatus</i> T. Scott (1894)
		<i>P. annandalei</i> (Sewell, 1919)
		<i>Pseudodiaptomus</i> sp. (Herrick, 1884)
	Augaptilidae	<i>Haloptilus</i> sp. Giesbrecht in Giesbrecht & Schmeil (1898)
Cyclopoida	Oithonidae	<i>Oithona nana</i> Giesbrecht (1892)
		<i>O. oculata</i> Farran (1913)
		<i>O. setigera</i> Dana (1849)

---

---

				<i>O. similis</i> Claus (1866)
				<i>O. simplex</i> Farran (1913)
				<i>O. tenuis</i> Rosendorn (1917)
				<i>O. rigida</i> Giesbrecht (1896)
				<i>O. brevicornis</i> Giesbrecht (1891)
				<i>O. spinirostris</i> Claus (1863)
				<i>Oithona</i> sp. Baird (1843)
	Harpacticoida		Miraciidae	<i>Miracia efferata</i> Dana (1849)
				<i>Macrosetella gracilis</i> Dana (1848)
				<i>M. oculata</i> Sars G.O. (1916)
			Ectinosomatidae	<i>Microsetella norvegica</i> Boeck (1864)
				<i>M. rosea</i> Dana (1848)
			Clytemnestridae	<i>Clytemnestra scutellata</i> Dana (1848)
			Euterpinidae	<i>Euterpina acutifrons</i> Dana (1848)
			Longipediidae	<i>Longipedia weberi</i> Scott A. (1909)
	Poecilostomatoida		Oncaeidae	<i>Oncaea confera</i> Giesbrecht (1891)
				<i>O. media</i> Giesbrecht (1891)
				<i>O. mediterranea</i> Claus (1863)
				<i>O. venusta</i> Philippi (1843)
				<i>Oncaea</i> sp. Philippi (1843)
			Sapphirinidae	<i>Copilia quadrata</i> Dana (1849)
				<i>Sapphirina auronitens</i> Claus (1863)
				<i>S. maculosa</i> Giesbrecht (1892)
				<i>S. ovatolanceolata</i> Dana (1849)
				<i>Sapphirina</i> sp. Thompson J. (1829)
			Corycaeidae	<i>Corycaeus agilis</i> Dana (1849)
				<i>C. andrewsi</i> Farran (1911)
				<i>C. catus</i> F. Dahl (1894)
				<i>C. danae</i> Giesbrecht (1891)
				<i>C. erythraeus</i> Cleve (1904)
				<i>C. lautus</i> Dana (1849)
				<i>C. longistylis</i> Dana (1849)
				<i>C. ovalis</i> Claus (1863)
				<i>C. robustus</i> Giesbrecht (1891)
				<i>C. speciosus</i> Dana (1849)
				<i>Corycaeus</i> sp. Dana (1845)
				<i>Farranula carinata</i> Giesbrecht (1891)
				<i>F. concinna</i> Dana (1849)
				<i>F. curta</i> Farran (1911)
				<i>F. gibbula</i> Giesbrecht (1891)
				<i>F. gracilis</i> Dana (1849)
Arthropoda:				
Ostracoda	Myodocopa	Myodocopida	Cypridinidae	<i>Macrocypridina castanea</i> Brady (1897)
		Halocyprida	Halocyprididae	<i>Conchoecia elegans</i> Sars (1865)
				<i>Euconchoecia chierchiaie</i> G. W. Müller (1890)
Arthropoda:				
Malacostraca	Eumalacostraca	Mysida	Mysidae	<i>Mesopodopsis orientalis</i> W. Tattersall (1908)
		Amphipoda	Leucothoidae	<i>Leucothoe spinicarpa</i> Abildgaard (1789)

---



		Hyperiididae	<i>Hyperia</i> sp. Latreille (1823)
		Caprellidae	<i>Caprella</i> sp. Lamarck (1801)
			<i>Metacaprella</i> sp. Mayer (1903)
		Talitridae	<i>Orchestoidea</i> sp. Nicolet (1849)
	Cumacea	Nannastacidae	<i>Campylaspis costata</i> Sars (1865)
	Euphausiacea	Euphausiidae	<i>Euphausia tenera</i> Hansen (1905)
			<i>Euphausia</i> sp. Dana (1850)
	Decapoda	Luciferidae	<i>Lucifer hanseni</i> Nobili (1905)
		Sergestidae	<i>Sergestes</i> sp. H. Milne Edwards (1830)
Chaetognatha:			
Sagittioidea	Aphragmophora	Sagittidae	<i>Pseudosagitta maxima</i> Conant (1896)
			<i>Sagitta bedoti</i> Beraneck (1895)
			<i>S. bipunctata</i> Quoy & Gaimard (1828)
			<i>S. enflata</i> Grassi (1881)
			<i>Sagitta</i> sp. Quoy & Gaimard (1827)
			<i>Adinosagitta bedfordii</i> Doncaster (1903)
Chordata:			
Appendicularia	Copelata	Oikopleuridae	<i>Oikopleura dioica</i> Fol (1872)
			<i>O. parva</i> Lohmann (1896)
		Fritillariidae	<i>Fritillaria</i> sp. Fol (1872)
Chordata:			
Thaliacea	Doliolida	Doliolidae	<i>Doliolum</i> sp. Quoy & Gaimard (1834)
	Salpida	Salpidae	<i>Salpa fusiformis</i> Cuvier (1804)

*Clytemnestra scutellata*, *Euterpina acutifrons*, *Longipedia weberi* of the harpacticoid were encountered as common species in this locality, only two species viz. *Euterpina acutifrons* and *Longipedia weberi* were more frequent. Three families viz. Oncaeidae, Sapphirinidae and Corycaeidae have represented the poecilostomatoida in which Corycaeidae with species *Corycaeus agilis*, *Corycaeus andrewsi*, *C. catus*, *Corycaeus sp.* and *Farranula gibbula* were dominant. Many times *Oncaea venusta* has also occurred quite frequently and that could be due to its diverse feeding habit and respiratory adaptation as opined earlier<sup>35</sup>. Seasonal variation in species dominance was observed in this part of Bay of Bengal. During premonsoon and monsoon season, *Paracalanus aculeatus* and *Acrocalanus longicornis* proved to be the most dominant species. Other species viz. *Oithona* sp., *Miracia efferata* and *Acrocalanus gibber* were dominant during premonsoon, while *Acartia southwelli*, *Acartia erythraea* and *Oithona* sp. were dominant during monsoon season. Faunal composition of copepod remained significantly different during post-monsoon period than premonsoon and monsoon season. The most dominant species were *Acrocalanus gibber*, *Centropages tenuiremis*, *Paracalanus parvus*, *Acrocalanus gracilis* and *Acartia erythraea*. Dominance of low saline species of genus *Paracalanus*, *Acartia* and

*Acrocalanus* in most part of the year indicate the estuarine influence in the study area.

The other crustacean fauna of the present study mainly belonged to three classes namely Malacostraca (mysids, cumaceans, euphausiids, amphipods and decapods), Ostracoda (ostracods) and Branchiopoda (cladocerans). In total 2 species of Cladocerans viz. *Evadne tergestina* and *Penilia avirostris* were encountered. They were more abundant (302 org./ 10m<sup>3</sup>) during POM season compared to other seasons. *Evadne tergestina* was commonly seen in plankton collections of POM period. Ostracoda population comprised of myodocopida mainly *Macrocypridina castanea* and halocyprida consisting of *Conchoecia elegans* and *Euconchoecia chierchiae*. The ostracoda population was represented by 3 species viz., *Macrocypridina castanea*, *Conchoecia elegans* and *Euconchoecia chierchiae* as against only one species i.e. *Philomedes* sp earlier<sup>17</sup>. Malacostraca comprised of 11 species belonging to 5 orders. Out of these 11 species, *Lucifer hanseni* occurred all year round and exhibited peak dominance during PRM period. This could be due to the fact that the Bay of Bengal remains more productive in this season and that supports for high Decapoda density.

The Protozoa population was represented by one species *Acanthometron* sp. of class Acantharia that occurred only in May and 15

species belonging to 8 families of class Ciliophora. Ciliophora population comprised of *Leptotintinnus nordqvisti*, *Tintinnopsis beroidea*, *Tintinnopsis butschlii*, *Tintinnopsis cylindrica*, *Tintinnopsis mortensenii*, *Tintinnopsis tocaninensis*, *Tintinnopsis tubulosa*, *Tintinnopsis uruguayensis*, *Codonellopsis ostenfeldi*, *Dictyocysta seshaiya*, *Metacylis jorgenseni*, *Rhabdonella* sp., *Amphorellopsis* sp., *Eutintinnus tenue*, *Favella philippinensis*. They were present during March, June, July, August, October,

Table 4. Check list of zooplankton (Meroplankton) encountered in the coastal waters of south Odisha coast, off Rushikulya estuary during 2010-11

#### Meroplankton

Actinula larva of Anthomedusae
Alima larva of Squilla
Brachiopod larvae
Brachyuran zoea larvae
Bryozoan cyphonautes larvae
Caridean larvae
Cypris larvae (Barnacle)
Cirripede nauplii (Barnacle)
Copepod nauplii
Gastropod veliger
Bivalve veliger
Isopod larvae
Larvae of euphausiids
Megalopa larvae of Brachyuran crab
Megalopa larvae of Pagurid crab
Planula larva of obelia
Polychaete larva
Protozoa of Lucifer
Zoea larva of porcelain crab
Echinoderm larvae
Ascidians larvae
Fish egg
Fish larvae

December 2010 and January 2011. The tintinnid population of this study resembled those in Bahuda estuary<sup>2</sup>, Vellar-Coleroon estuary<sup>36</sup> and coastal regions of Sundarban mangrove wetland<sup>37</sup>. Foraminifera were represented by 5 species *viz.* *Globigerina bulloides*, *Globigerina rubescens*, *Globigerina* sp., *Asterorotalia trispinosa* and *Quinqueloculina* sp. and were common during March - May and November - February.

Hydrozoa of phylum Cnidaria were represented by 18 species belonging to two classes namely hydroidomedusae and Siphonophorae. They were present throughout the

year except in September. The peak period of Siphonophorae occurrence was observed during February and March. *Diphyes dispar* appeared as the most common siphonophorae in this area.

Ctenophora was represented by two families i.e. Pleurobrachiidae and Beroidea with a single species *viz.* *Pleurobrachia pileus* and *Beroe* sp. respectively. *P. pileus* was commonly seen during MON and POM and was completely absent during PRM season. *Beroe* sp. on the other hand had occurred only during POM and its population size was poor.

Phylum Mollusca was represented by Gastropoda comprising of 6 species namely *Atlanta* sp., *Janthina* sp., *Limacina bulimoides*, *Limacina inflata*, *Cresis acicula* and *Hyalocylis striata*. Among these species *Cresis acicula* was dominant which occurred in greater part of the year. Its presence was more conspicuous in March and December.

The Chaetognatha population was represented by six species *namely* *Pseudosagitta maxima*, *Sagitta bedoti*, *Sagitta bipunctata*, *Sagitta enflata*, *Sagitta* sp. and *Adinosagitta bedfordii* under a single family Sagittidae. They were more abundant during the PRM and POM months. *Sagitta enflata* was the most dominant species during our study which is comparable to many observations<sup>38</sup>.

Planktonic chordates of the region were represented by 3 species of appendicularians *viz.* *Oikopleura dioica*, *Oikopleura parva* and *Fritillaria* sp. and one species each of Dollioloids (*Dolliolum* sp.) and salps (*Salpa fushiformis*). *O. dioica* however remained as the most common species among the appendicularians with high abundance during March compared to other months.

Meroplankton was represented by 23 forms during the study. They were more abundant during the PRM period compared to other season. Among these 23 forms, bivalve veliger, brachyuran zoea larva, caridean larva, cirripede nauplii, copepod nauplii, fish egg & larvae, gastropod veliger, polychaete larva, protozoa of lucifer were frequently observed throughout the year.

Species diversity of plankton community exhibited well marked fluctuations (Table 7). The Shannon Wiener Diversity index ( $H'$  (loge)) was higher in February (3.41), while the Margalef's species richness ( $d$ ) (8.32) & Simpson's dominance ( $\lambda$ ) (0.22) in January, and Pielou's evenness ( $J'$ ) in February (0.78). Higher

Simpson's Dominance ( $\lambda$ ) (0.22) as well as lower Shannon Wiener Diversity index (2.02) during January might be due to the dominance of species like *Paracalanus parvus*, *Farranula gibbula*, *Corycaeus erythraeus*. Seasonally the Shannon diversity and evenness of zooplankton community were generally higher in monsoon while richness and dominance during POM period.

Table 5 and Fig. 2 represents the population density and percentage composition of different groups of zooplankton respectively. The population density of zooplankton exhibited wide range of spatial and seasonal variations. The average density varied between 1077 org./10m<sup>3</sup> (station R-3 in MON) and 23060 org./10m<sup>3</sup> (station R-1 in PRM). The seasonally pooled




Fig.2. Relative abundance of zooplankton community during 2010-11

Table 5. Monthly variation in population size of different zooplankton groups during 2010-11

Major Groups	Mar	Apr	May	Jun	PRM	Jul	Aug	Sep	Oct	MON	Nov	Dec	Jan	Feb	POM
Acantharia	0	0	37	0	9	0	0	0	0	0	0	0	0	0	0
Ciliophora	29	0	0	6	9	1	2	0	38	10	0	5	0	0	1
Foraminifera	35	6	37	0	19	0	0	0	0	0	3	87	3	12	26
Hydrozoa	236	99	51	21	102	1	6	0	12	5	43	26	7	73	37
Gastropoda	161	11	0	29	50	3	2	0	4	2	6	403	2	77	122
Cladocera	141	0	0	27	42	72	1	107	92	68	411	12	725	58	302
Copepoda	11946	7731	9677	2979	8083	1141	1062	3067	2438	1927	4799	7627	2837	2996	4565
Ostracoda	37	0	0	58	24	16	9	0	21	12	0	2	0	5	2
Malacostraca	232	65	69	74	110	45	23	157	41	67	89	137	0	17	61
Chaetognatha	0	71	40	25	34	6	14	0	21	10	29	220	35	51	84
Chordata	520	17	69	13	155	1	1	0	6	2	33	6	25	8	18
Meroplankton	7342	1554	2872	320	3022	166	144	423	402	284	277	1005	187	557	507

average zooplankton density showed that the rank order distribution of population size in different seasons (Table 1) are PRM (11659 org./10m<sup>3</sup>) > POM (5725 org./10m<sup>3</sup>) > MON (2387 org./10m<sup>3</sup>). The higher PRM population density of zooplankton of the present study coincides with

many earlier reports from the coastal waters of Indian seas<sup>34, 39</sup> and other literature cited thereon. Station-wise population density of different groups of zooplankton during PRM period have shown that the rank order distribution of different groups is copepod > larvae > malacostraca >

others in station R-1 and R-2, whereas in station R-3 and R-5 the population density although followed the same sequence, appendicularia has occupied the 3rd place in place of malacostraca followed by others. At station R-4, the rank order distribution stands as copepod > larvae > hydrozoa. Similar to PRM, the rank order distribution of population density of MON season showed that, copepods, larvae and malacostraca occupied the 1st, 2nd and 3rd position in order of their dominance at stations R-1, R-2 and R-5, while at stations R-3 and R-4, branchiopoda was in third position in place of malacostraca. During post MON months, the zooplankton population too exhibited well defined spatial variation. Like PRM and MON, copepoda remained as the most dominant group. It was followed by larval forms at stations R-2, R-3 and R-5 and branchiopoda and gastropoda as the 2nd dominant group at stations R-1 and R-4 respectively. Group wise contributions of major taxa are given here under.

The population density of copepods showed well marked spatial and temporal variation (Table 5). Average numerical abundance of copepods was maximum in March (11946 org./10m<sup>3</sup>) and minimum in August (1062 org./10m<sup>3</sup>). Seasonal pooled average population density of copepod was higher during PRM (8083 org./10m<sup>3</sup>) followed by POM (4565 org./10m<sup>3</sup>) and MON (1927 org./10m<sup>3</sup>). Higher population density of copepod during PRM suggested higher secondary productivity. Similar situation was also reported earlier along the Odisha coast<sup>2,11-13</sup>. The occurrence of higher values of copepod density in the zooplankton community corroborates many earlier findings<sup>33,35,40-42</sup>. During the present study, contribution of Copepod to the total zooplankton population ranged from 58-84% (Fig.2). Many workers have also reported almost similar contribution of copepods ranging from 51.82-91.25% in Rushikulya estuary<sup>13</sup>, 57.94-89.98% in Bahuda estuary<sup>2</sup>, in Chilika lake 42.58-69.69% by Devasundaram and Roy<sup>10</sup>, 67.84-94.99% by Naik *et al*<sup>12</sup> & and 53.5-67.9% in the coastal waters of Bay of Bengal off the Rushikulya estuary<sup>17</sup>. Calanoid copepods contributed 64.0%, 73.7% and 82.2% to total copepod during PRM, MON and POM season respectively. Cyclopoids contributed to about 17.6%, 17.7%, 2.5%, where as harpacticoida contributed to 11.5%, 2.5%, 2.7% corresponding to the PRM, MON and POM seasons respectively. Poecilostomatoida contributed only to about 6.9%, 6.1% and 12.6% during PRM, MON and POM periods. Calanoid dominance as observed during the

present study is common phenomenon in Indian coastal waters.

Monthly analysis showed that Cladoceran was more abundant during January (725 org./10m<sup>3</sup>), November (avg. 411 org./10m<sup>3</sup>) and March (avg.141 org./10m<sup>3</sup>) which coincided with the observations of Della and Venugopal<sup>43</sup>. They were abundant during POM (302 org./10m<sup>3</sup>) and MON (68 org./10m<sup>3</sup>) period as reported earlier by Naomi *et al*<sup>44</sup>. Among cladocerans, *Evadne tergestina* was more abundant and with higher average population density of 298 org./10m<sup>3</sup> during POM season followed by MON (51 org./10m<sup>3</sup>) and PRM (4 org./10m<sup>3</sup>). The highest population density (58 org./10m<sup>3</sup>) of ostracods was encountered in June. Seasonally pooled density of this group varied between 2 (POM) & 24 (PRM) org./10m<sup>3</sup>. The seasonal average population density of malacostraca represented by 5 forms namely mysids, cumaceans, euphausiids, amphipods & decapods ranged between 61-110 org./10m<sup>3</sup> with maximum and minimum density observed during PRM and POM respectively.

The protozoa *Acanthometron* sp. occurred only in the month of May with population density of 37 org./10m<sup>3</sup>. Highest population density (38 org./10m<sup>3</sup>) of tintinnids was encountered in October followed by March (29 org./10m<sup>3</sup>). In other months it was either scarce or absent. Low population size of tintinnids could be due to the use of large mesh size (120µm) plankton net as was reported earlier<sup>45</sup>.

The population density of foraminifera ranged from 3 - 87 org./10m<sup>3</sup> and were totally absent during the MON season. Pooled average concentration of this group varied from 19 (PRM) - 26 (POM) org./10m<sup>3</sup> (Table 5). Presence of benthic foraminifera in the plankton collection could be ascribed to their addition from the bottom as a result of upward movement in water column as was reported earlier<sup>46</sup>.

Group Hydrozoa was encountered during all the months except in September. Their population density ranged from 1-236 org./10m<sup>3</sup>. Maximum seasonal average density of 102 org./10m<sup>3</sup> was noticed during PRM in which hydroidomedusae were 89 org./10m<sup>3</sup>. Such higher population density during PRM was also reported from Indian coasts<sup>47</sup>. The peak periods of siphonophores abundance in March and February is in coincidence with the earlier observation in Cochin Backwater<sup>48</sup>. Among the 10 species of siphonophores, *Diphyes dispar* remained as the dominant species contributing 78.4% in PRM,

55.4% in MON and 82.0% in POM of total siphonophore population.

Gastropoda had occurred throughout the year except in May and September. They were more abundant (403 org./10m<sup>3</sup>) during December (Table 5). Their population size ranged from 2 - 403 org./10m<sup>3</sup>. Seasonally pooled average density varied from 2(MON) - 122 (POM) org./10m<sup>3</sup>. Of all the species, *Cresis acicula* appeared during greater part of the year and were more conspicuous in December (402 org./10m<sup>3</sup>) which resembles with the observation in inshore waters of Karwar<sup>49</sup>.

The population size of chaetognaths ranged from 6 - 220 org./10m<sup>3</sup>. They were encountered in good numbers in December. Seasonally pooled average values showed that it remained higher during POM season (84 org./10m<sup>3</sup>) followed by PRM (34 org./10m<sup>3</sup>) and MON (10 org./10m<sup>3</sup>) in order. High abundance of chaetognaths during POM was also reported by in the coastal water of Adubidri<sup>39</sup>. Compared with the other groups, Ctenophora appeared with very low percentage with a maximum density of 2 org./10m<sup>3</sup>. The population density of Chordates represented by two major classes namely appendicularia and thaliacea ranged from 1-520 org./10m<sup>3</sup> in which *Oikopleura dioica* was more abundant. *Doliolum* sp and *Salpa fusiformis* belonging to thaliacea were found but was in very low percentages.

In total, 23 different larval forms of invertebrates were observed during the study period (Table 4). Larval population showed their maximum density during March (7342

org./10m<sup>3</sup>). Average PRM density was 3022 org./10m<sup>3</sup> followed by POM (507 org./10m<sup>3</sup>) and MON (284 org./10m<sup>3</sup>). Gastropod veliger, Caridean larvae and fish eggs were dominant in the larval population of PRM, MON and POM respectively. Gastropod veligers were more abundant in March compared to other months. It agrees with the observations made in the inshore waters of Karwar<sup>49</sup>. All the crustacean larvae together contributed 3.8% in POM - 9.0 % in PRM of the total zooplankton density which was high as compared to the earlier report<sup>17</sup>. The other larvae together contributed to about 4.2% in MON to 17.0% during PRM of the total zooplankton density.

Zooplankton density showed positive correlation with DO ( $r=0.546$ ,  $p<0.05$ ), nitrate ( $r=0.672$ ,  $p<0.01$ ) and chlorophyll *a* ( $r=0.857$ ,  $p<0.01$ ) concentration (Table 6). The positive correlation values between zooplankton density and chlorophyll *a* suggested the coexistence of zooplankton community and phytoplankton. Normally the zooplankton density exhibits inverse relationship with phytoplankton. However positive correlations between these two groups are also not very rare as opined by Prasad<sup>50</sup>. Many researchers<sup>40,42</sup> have reported direct relationship between salinity and zooplankton population density. In the present study, however, no definite correlation between these two parameters was observed.

Non-metric multidimensional scaling (MDS) ordinations based on Bray-Curtis<sup>51</sup> similarities of species abundance data were produced to provide

Table 6. Correlation matrix among different physicochemical and biological parameters

	WT	pH	DO	Salinity	NO <sub>2</sub>	NO <sub>3</sub>	PO <sub>4</sub>	SiO <sub>4</sub>	Chl- <i>a</i>	Zoo
WT	1									
pH	.184	1								
DO	-0.155	-0.490	1							
Salinity	.629*	.687**	-.590*	1						
NO <sub>2</sub>	-0.218	-.525*	.279	-0.340	1					
NO <sub>3</sub>	-0.455	-0.259	.240	-.552*	-0.078	1				
PO <sub>4</sub>	-0.419	-.640*	.860**	-.712**	.345	.419	1			
SiO <sub>4</sub>	-.747**	-0.166	.395	-.583*	.073	.551*	.572*	1		
Chl- <i>a</i>	.080	-.544*	.356	-0.342	.042	.522*	.359	.057	1	
Zoo	.099	-0.379	.546*	-0.395	-0.086	.672**	.448	.155	.857**	1

\*Correlation is significant 0.05 level, \*\*Correlation is significant 0.01 level

a visual representation in a two-dimensional plot of the relative similarities in zooplankton community species composition and abundance at different sampling sites as well as on different

sampling occasions. This analysis formed different groups by taking into account of similar species composition (Fig. 3). From the MDS ordination (Fig. 3) it is clear that species composition was unequal during different months as there is no similarity found more than 40%

level. September, March, June and January were individually grouped proving their unique species composition, whereas cluster between some months were found irrespective of seasons. This type of poor similarity index could be attributed to the patchy distribution of zooplankton in this area as observed by Omori & Hamner<sup>52</sup>. From these results it could be inferred that there is a spatial and temporal heterogeneity in species composition and distribution of zooplankton in this part of the Bay of Bengal.

Table 7. Monthly variations of Univariate diversity indices

Sample	d	J'	H'(loge)	Lambda
Mar	6.76	0.74	3.22	0.08
Apr	5.11	0.65	2.64	0.13
May	4.16	0.70	2.70	0.10
Jun	6.04	0.66	2.69	0.14
Jul	7.20	0.77	3.21	0.06
Aug	5.26	0.70	2.71	0.14
Sep	4.68	0.71	2.74	0.12
Oct	5.39	0.69	2.74	0.14
Nov	6.34	0.66	2.76	0.14
Dec	7.15	0.64	2.80	0.11
Jan	8.32	0.46	2.02	0.22
Feb	8.11	0.78	3.41	0.05




Fig.3. Non-parametric multi-dimensional scaling (MDS) ordination of the square-root transformed zooplankton community of each month

### Conclusion

In the present study effort was made to explore the zooplankton diversity as well as to determine current status of zooplankton composition and its population size variation in coastal waters of the Bay of Bengal, off the Rushikulya estuary. It has been found that the faunal composition of zooplankton remained significantly diverse. Copepods have emerged as the most dominant group contributing >50% of

total population. Occurrence of low saline copepod species signified estuarine influence on the distribution of plankton community. It further revealed that zooplankton fauna of the region is susceptible to change under the influence of different environmental parameters such as salinity, chlorophyll *a* and nutrients of the ambient medium, rather than salinity resulting heterogeneity in species composition, population size and abundance of zooplankton.

### Acknowledgement

Authors wish to thank the Head of the Department of Marine Sciences for providing required facilities and encouragements. The facilities of the INCOIS sponsored SATCORE project was availed for field collection and analysis work. One of the authors (BKS) wishes to thank DST, Govt. of India for providing fellowship.

### Reference

- Roman, M., Smith, S., Wishner, K., Zhang, X. & Gowing, M., Mesozooplankton production and grazing in the Arabian Sea, Deep-Sea Res. II, 47(2000) 1423-1450.
- Mishra, S. & Panigrahy, R. C., Zooplankton ecology of the Bahuda estuary (Orissa), east coast of India, Indian J. Mar. Sci., 28(1999) 297-301.
- Rao, T.S.S. & Ganapati, P.N., Studies on Chaetognatha in Indian seas. Part-III. Andhra Univ. Mem. Oceanogr., 2(1958) 147-163.
- Bonnet, D. & Frid, C. S., Copepod species considered as indicators of water mass influence and changes: results from a Northumberland coastal station, ICES. J. Mar. Sc., 61(2004) 485-491.
- Sewell, R. B. S., Notes on plankton from the Chilka Lake, Rec. Indian Mus., 9(1913) 338-340.
- Madhu, N. V., Jyothibabu, R., Balachandran, K. K., Honey, U. K., Martin, G. D., Vijay, J. G., Shiyas, C. A., Gupta, G. V. M. & Achuthankutty, C. T., Monsoonal impact on planktonic standing stock and abundance in a tropical estuary (Cochin backwaters, India), Estuar. Coast. Shelf Sci., 73(1-2) (2007) 54-64.
- Dalal, S. G. & Goswami, S. C., Temporal and ephemeral variations in copepod community in the estuaries of Mandovi and Zuari west coast of India, J. Plank. Res. 23(1) (2001) 19-26.
- Santhanam, P. & Perumal, P., Diversity of zooplankton in Parangipettai coastal waters, southeast coast of India, J. Mar. Biol. Assoc. India, 45(2003)144-151.
- Sarkar, S. K., Singh, B. N. & Choudhury, A., Composition and variations in the abundance of zooplankton in the Hoogly estuary, West Bengal, India, Proc. Indian Acad. Sci., 95(2) (1986) 125-134.
- Devasundaram M P & Roy J C, A preliminary study of the plankton of the Chilka Lake for the years 1950 and 1951, Symposium on Marine and Fresh Water Plankton in the Indo-Pacific, IPFC Publication, 1954.
- Patnaik, S., Observations on the seasonal fluctuations of plankton in the Chilka Lake, Indian J. Fish., 20(1973) 43-55.
- Naik, S., Panigrahy, R. C. & Mohapatro, A., Spatio-temporal distribution of zooplankton in Chilka lake- A

- Ramsar site on the Indian east coast, Indian J. Sci. Tech., 3(2008) 1-5.
13. Gouda, R. & Panigrahy, R. C., Zooplankton ecology of the Rushikulya estuary, East coast of India, J. Aquacult. Tropics, 10(1995) 201-211.
  14. Mishra S. & Panigrahy, R. C., Copepods of Bahuda estuary (Orissa), east coast of India, Indian J. Mar. Sci., 25(1996) 98-102.
  15. Ramaiah, N., Chatterji, A. & Madhupratap, M., A study on the zooplankton of the Burhabalanga estuary, Orissa coast, Proc. Indian National Sci. Aca., 62(1996) 1-4.
  16. Srichandan, S., Panda, C. R. & Rout, N.C., Seasonal distribution of zooplankton in Mahanadi Estuary (Odisha), East Coast of India: A Taxonomical Approach, Int. J. Zool. Res., 9(1) (2013) 17-31.
  17. Sahu, G., Mohanty, A. K., Singhasamanta, B., Mahapatra, D., Panigrahy, R. C., Satpathy, K. K. & Sahu, B.K., Zooplankton Diversity in the Nearshore Waters of Bay of Bengal, Off Rushikulya Estuary, The IUP J. Environ. Sci., 4(2010) 61-85.
  18. Mahapatra, T.R. & Padhy, S.N., Nutrients and physicochemical characterisation of Rushikulya estuary during PRM season, J. Env. Poll., 8 (1) (2001) 61-65.
  19. Grasshoff K, Ehrhardt M & Kremling K, Methods of seawater analysis, 3rd.edition, (Verlag Chemie GmbH, Weinheim) 1999, pp. 600.
  20. Parsons T R, Maita Y & Lalli C M, A Manual of Chemical and Biological. Methods for Seawater Analysis (Pergamon Press, New York) 1984, pp. 173.
  21. Kasturirangan L R, A key for the identification of the more common planktonic Copepoda of Indian coastal waters, Indian National Committee on Oceanic Research, Council of Scientific and Industrial Research, New Delhi publication No. 2, 1963.
  22. Newell G E & Newell R C, *Marine Plankton- A Practical Guide* (Hutchinson Educational Ltd., London) 1977, pp.244.
  23. Conway D V P, White R G, Hugues-Dit-Ciles J, Gallienne, C P & Robins D B., *Guide to the coastal and surface zooplankton of the South-Western Indian Ocean*. Publication No.15, (Marine Biological Association of the United Kingdom) 2003, pp.354.
  24. Clarke K R & Gorley R N, *PRIMER v5 : User manual/tutorial, PRIMER-E*, (Plymouth UK), 2001, pp. 91.
  25. Clarke K R & Warwick RM, Change in marine communities: an approach to statistical analysis and interpretation, 2nd edition (PRIMER-E, Plymouth) 2001.
  26. Kannan, R. & Kannan, L., Physico-Chemical characteristics of seaweed beds of the Palk Bay, Southeast coast of India, Indian J. Mar. Sci., 25(1996)358-362.
  27. Satpathy, K. K., Seasonal distribution of nutrients in the Coastal waters of Kalpakkam, East coast of India, Indian J. Mar. Sci., 25(1996) 221-224.
  28. Subramanian, B. & Mahadevan, A., Seasonal and diurnal variation of hydrobiological characters of coastal waters of Chennai, Bay of Bengal, Indian J. Mar. Sci., 28(1999) 429-433.
  29. Satpathy, K. K., Mohanty, A K., Sahu, G., Prasad, M. V.R., Venkatesan, R., Natesan, U. & Rajan, M., On the occurrence of *Trichodesmium erythraeum*(Ehr.) bloom in the coastal waters of Kalpakkam, east coast of India, Indian J. Sci. Tech, 1(2007) 1-11.
  30. Govindasamy, C., Kannan, L. & Azariah, J., Seasonal variation in physico-chemical properties and primary production in the coastal water biotopes of Coromandel coast, India, J. Environ. Biol., 21(2000) 1-7.
  31. Choudhury, S.B. & Panigrahy, R.C., Seasonal distribution and behavior of nutrients in the creek and coastal waters of Gopalpur, east coast of India, Mahasagar, 24(1991) 81-88.
  32. Ramaiah, N. & Nair, V., Distribution and abundance of copepods in the pollution gradient zones of Bombay harbour- Thane creek- Basin creek, west coast of India, Indian J. Mar. Sci., 26(1997) 20-25.
  33. Rakesh, M., Raman, A.V. & Sudarsan, D., Discriminating zooplankton assemblages in neritic and oceanic waters: A case for the northeast coast of India, Bay of Bengal, Mar. Environ. Res., 61 (2006) 93-109.
  34. Santhosh Kumar, C. & Perumal, P., Hydrobiological Investigations in Ayyampattinam Coast (Southeast Coast of India) with special reference to zooplankton, Asian J. Biol. Sci., 4(1) (2011) 25-34.
  35. Fernandes, V. & Ramaiah, N., Mesozooplankton community in the Bay of Bengal (India): spatial variability during the summer monsoon, Aquat. Ecol., 43 (4) (2009) 951-963.
  36. Godhantaraman, N., Seasonal variations in species composition, abundance, biomass and estimated production rates of tintinnids at tropical estuarine and mangrove waters, Parangipettai, southeast coast of India, J. Mar. Sys., 36 (2002) 161- 171.
  37. Biswas S. N., Godhantaraman N., Rakshit, D. & Sarkar, S.K., Community composition, abundance, biomass and production rates of Tintinnids (Ciliata: protozoa) in the coastal regions of Sundarban Mangrove Wetland, India, Indian J. Geo. Mar. Sci., 42 (2) (2013) 163-173.
  38. George, R. M., Thomas, P. A., Jasmine, S., Nair, K. R. & Vasanthakumar, R., Observations on the distribution and seasonal fluctuations of Chaetognaths off Vizhinjam, Southwest Coast of India, J. Mar. Biol. Ass. India, 40 (1 & 2) (1998) 6 - 10.
  39. Resmi, S., Reddy, R.V., Venkatesha Moorthy, K.S. & Chethan, N., Zooplankton dynamics in the coastal waters of adubidri, Karnataka, Indian J. Geo-Mar. Sci., 40(1) (2011) 134-141.
  40. Damotharan, P., Perumal, N.V., Arumugam, M., Perumal, P., Vijayalakshmi, S., & Balasubramanian, T., Studies on zooplankton Ecology from Kodiakkarai(Point Calimer Coastal Waters (South East Coast of India), Res. J. Biol. Sci., 5(2) (2010)187-198.
  41. Fernandes, V., The effect of semi-permanent eddies on the distribution of mesozooplankton in the central Bay of Bengal, J. Mar. Res., 66(4) (2008) 465-488.
  42. Jagadeesan, L., Jyothibabu, R., Anjusha, A., Mohan Arya, P., Madhu, N.V., Muraleedharan, K.R. & Sudheesh, K., Ocean currents structuring the mesozooplankton in the Gulf of Mannar and the Palk Bay, southeast coast of India, Progr. Oceanogr., (2013) 1-22.
  43. Della, C. N. & Venugopal, P., Distribution of marine cladocerans in the Indian Ocean, Mar. Biol., 16(1972)132-138.
  44. Naomi T S, Antony G & Mathew K J, Studies on the distribution of cladocera in the eastern Arabian sea and the Bay of Bengal, in: *Proceeding of First workshop on Scientific Results on FORV Sagar Sampada*, 1990, pp.85-93.
  45. Sieburth, JMc., Smetacek, V. & Lenx, J., Pelagic ecosystem structure: Heterotrophic compartments of the plankton and their relationship to plankton size fractions, Limnol. & Oceanogr., 23 (1978) 1256-1263.
  46. Murray, J. W., Significance of benthic foraminiferids in plankton samples, J. Paleol., 39(1) (1965) 156-157.
  47. Santhakumari, V. & Nair, V. R., Distribution of hydromedusae from the exclusive economic zone of the west and east coast of India, Indian J. Mar. Sci., 28(1999) 150-157.

48. Rengarajan, K., On the occurrence of siphonophores in the Cochin Backwater, *J. Mar. Biol. Assoc. India*, 16 (1) (1974) 280-285.
49. Naomi, T .S., On the zooplankton of the inshore waters of Karwar during 1980-81, *Indian J. Fish.*, 33 (3) (1986) 336-346.
50. Prasad, R. R., Further studies on the plankton of the inshore waters off Mandapam, *Indian J. Fish.*, 3(1) (1956) 1-42.
51. Bray, J. R. & Curtis, J. T., An ordination of the upland forest communities of Southern Wisconsin, *Ecol. Monogr.*, 27 (1957) 325-349.
52. Omori, M. & Hamner, W.M., Patchy distribution of zooplankton: behavior, population assessment and sampling problems, *Mar. Biol.*, 72(2) (1982) 193-20